

# The quartic equation: invariants and Euler's solution revealed<sup>1</sup>

RWD Nickalls<sup>2</sup>

*The Mathematical Gazette* (2009); vol. 93 (March, No 526), p. 66–75  
<http://www.nickalls.org/dick/papers/maths/quartic2009.pdf>

## 1 Introduction

The central role of the resolvent cubic in the solution of the quartic was first appreciated by Leonard Euler (1707–1783). Euler's quartic solution first appeared as a brief section (§5) in a paper on roots of equations [1, 2], and was later expanded into a chapter entitled *Of a new method of resolving equations of the fourth degree* (§§773–783) in his *Elements of algebra* [3, 4].

Euler's quartic solution was an important advance, in which he showed that each of the roots of a reduced quartic can be represented as the sum of three square roots, say  $\pm\sqrt{r_1} \pm \sqrt{r_2} \pm \sqrt{r_3}$ , where the  $r_i$  ( $i = 1, 2, 3$ ) are the roots of a resolvent cubic. A quartic equation in  $x$  is said to be *reduced* if the coefficient of  $x^3$  is zero. This can always be achieved by a simple change of variable.

Motivated by the recent tercentenary of Euler's birth, this article describes the geometric basis underlying both the  $r_i$  and the sign of the product  $\sqrt{r_1 r_2 r_3}$ , these being two key aspects of Euler's solution. Finally, we reveal the beautiful dynamic between Euler's resolvent cubic and the quartic invariants  $G, H, I, J$  [5, 6, 7], and propose a new class of algebraic object.

## 2 Geometric basis for the $r_i$

A significant property of the reduced quartic equation is that the four roots can be completely defined using only three parameters. For example, let  $z_j$  ( $j = 1, 2, 3, 4$ ) be the roots (see Figure 1) of a reduced quartic equation,

$$Z(x) \equiv ax^4 + px^2 + qx + r = 0. \quad (1)$$

As the sum of the roots is zero (the coefficient of the cubic term is zero), it follows that we can define the points midway between  $z_1, z_2$  and  $z_3, z_4$  as  $\pm g$ . Let  $z_2 - z_1 = 2\alpha$  and  $z_4 - z_3 = 2\beta$ . The four roots can then be expressed as follows:

$$\begin{cases} z_1, z_2 = -g \pm \alpha, \\ z_3, z_4 = +g \pm \beta. \end{cases}$$

---

<sup>1</sup>This minor revision of the original article corrects typographic errors and incorporates some explanatory footnotes. The original published version will soon be available in the JSTOR archive.

<sup>2</sup>Department of Anaesthesia, Nottingham University Hospitals, City Hospital Campus, Nottingham, UK. email: [dick@nickalls.org](mailto:dick@nickalls.org) [[www.nickalls.org](http://www.nickalls.org)]




Figure 1:

Since specifying one pair of quartic roots necessarily defines the remaining pair, there are just three different ways of allocating the pairs of roots, each associated with its own  $g, \alpha, \beta$ , the inter-relationship between which lies at the heart of a remarkable symmetry which underpins the solution of the quartic. For example if, with no loss of generality, we let

$$\begin{cases} z_3 + z_4 = 2g_1, \\ z_3 + z_1 = 2g_2, \\ z_3 + z_2 = 2g_3, \end{cases} \quad (2)$$

then

$$\begin{aligned} 2(g_2 + g_3) &= 2z_3 + z_1 + z_2, \\ &= (z_1 + z_2 + z_3 + z_4) + z_3 - z_4, \\ &= z_3 - z_4 = -2\beta_1, \end{aligned}$$

and similarly

$$2(g_2 - g_3) = z_1 - z_2 = -2\alpha_1,$$

and hence

$$\begin{cases} \alpha_1 = -(g_2 - g_3), \\ \beta_1 = -(g_2 + g_3). \end{cases}$$

Thus the  $\alpha_k, \beta_k$  ( $k = 1, 2, 3$ ) are actually simple functions of the  $g_i$  ( $i \neq k$ ) such that each of the four roots  $z_j$  can be expressed as a function of the  $g_i$  alone, as follows:

$$\begin{cases} z_1 = -g_1 - \alpha_1 = -g_1 + (g_2 - g_3) = -g_1 + g_2 - g_3, \\ z_2 = -g_1 + \alpha_1 = -g_1 - (g_2 - g_3) = -g_1 - g_2 + g_3, \\ z_3 = +g_1 - \beta_1 = +g_1 + (g_2 + g_3) = +g_1 + g_2 + g_3, \\ z_4 = +g_1 + \beta_1 = +g_1 - (g_2 + g_3) = +g_1 - g_2 - g_3. \end{cases} \quad (3)$$

Thus Euler's  $r_i$  are the same as the  $g_i^2$ .

### 3 Euler's resolvent cubic

Using these observations we can reconstruct a given reduced quartic equation, say (1), which then leads to a resolvent cubic and hence to the solution. Let the roots of  $Z(x) = 0$  be  $-g \pm \alpha$  and  $g \pm \beta$  (Figure 1).

$$Z(x) \equiv \{x - (-g - \alpha)\}\{x - (-g + \alpha)\}\{x - (g - \beta)\}\{x - (g + \beta)\} = 0.$$

Expanding and letting  $A = g^2 - \alpha^2$  and  $B = g^2 - \beta^2$ , gives

$$x^4 + (-4g^2 + A + B)x^2 + (2g)(B - A)x + AB = 0.$$

We can eliminate  $\alpha, \beta$  by first equating coefficients with the monic form of (1) giving

$$\begin{cases} p/a &= -4g^2 + A + B, \\ q/a &= 2g(B - A), \\ r/a &= AB, \end{cases}$$

and then eliminating  $A$  and  $B$  (using the identity  $4AB = 2A \times 2B$ ), which generates a resolvent sextic in  $g$ , the roots of which are the six values  $\pm g_1, \pm g_2, \pm g_3$ . The substitution  $g^2 \mapsto x$  then generates Euler’s original resolvent cubic [1, 2, 3, 4]

$$R(x) \equiv x^3 + \frac{p}{2a}x^2 + \left(\frac{p^2 - 4ar}{16a^2}\right)x - \frac{q^2}{64a^2} = 0, \tag{4}$$

whose roots  $r_i$  are therefore  $g_1^2, g_2^2, g_3^2$ . The four roots of the reduced quartic  $Z(x) = 0$  are among the eight possible values of  $\pm\sqrt{r_1} \pm \sqrt{r_2} \pm \sqrt{r_3}$ ; but in order to determine which four they are we need a way of allocating the signs correctly.

Euler, using a monic quartic of the form  $x^4 - lx^2 - mx - n = 0$ , says he resolved the sign problem by noting that  $\sqrt{r_1 r_2 r_3} = m/8$ , as follows [3, § 773]:

... But it is to be observed, that the product ...  $\sqrt{r_1 r_2 r_3}$ , must be equal to  $m/8$ , and that if  $m/8$  be positive, the product of the terms  $\sqrt{r_1}, \sqrt{r_2}, \sqrt{r_3}$ , must likewise be positive;

Unfortunately Euler did not elaborate further on this, but the key to understanding the sign problem is not difficult to find, since from (2) we have

$$\begin{aligned} 8g_1g_2g_3 &= (z_3 + z_4)(z_3 + z_1)(z_3 + z_2), \\ &= z_3^3 + z_3^2(z_1 + z_2 + z_4) + z_3(z_2z_1 + z_2z_4 + z_1z_4) + z_4z_1z_2. \end{aligned}$$

Now  $z_1 + z_2 + z_4 = -z_3$  (since  $\Sigma z_j = 0$ ), hence

$$8g_1g_2g_3 = z_1z_2z_3 + z_2z_3z_4 + z_3z_4z_1 + z_4z_1z_2, \tag{5}$$

and so  $8g_1g_2g_3$  is actually one of the four elementary symmetric functions of the roots  $z_j$ . Its value is therefore equal to  $-1 \times$  the coefficient of the  $x$ -term of the monic form of the reduced quartic equation  $Z(x) = 0$ , and so we have

$$8\sqrt{r_1 r_2 r_3} = 8g_1g_2g_3 = -q/a, \tag{5a}$$

which is equivalent to Euler’s  $\sqrt{r_1 r_2 r_3} = m/8$ .

## 4 Geometric basis for the sign of $\sqrt{r_1 r_2 r_3}$

A useful way of ‘seeing’ the quartic algebra at work is to express the coefficients in terms of the key ‘visible’ parameters  $\varepsilon, y_{N_z}, y_{N_z'}$ , shown in Figure 2, as follows: Let  $F(X)$  be a quartic polynomial with real coefficients ( $a \neq 0$ )

$$F(X) \equiv aX^4 + bX^3 + cX^2 + dX + e, \tag{6}$$

with invariants [6, p. 76]

$$\begin{cases} G &= b^3 + 8a^2d - 4abc, \\ H &= 8ac - 3b^2, \\ I &= 12ae - 3bd + c^2, \\ J &= 72ace + 9bcd - 27ad^2 - 27eb^2 - 2c^3. \end{cases} \tag{7}$$




Figure 2:

The reduced quartic  $Z(x)$ , turning points  $(T_1, T_2, T_3)$ , points of inflection  $(I_1, I_2)$ , and first differential  $Z'(x)$ . The  $x$ -coordinates of the points of inflection are  $\pm\epsilon$ . The curves intersect the  $y$ -axis at points  $N_z$  and  $N_{z'}$ .

Let its reduced form  $Z(x)$  be generated by the translation  $x \mapsto X - X_{Nf}$ , where  $X_{Nf} = -b/(4a)$ . Using Taylor's theorem we have

$$Z(x) \equiv F(x + X_{Nf}) = ax^4 + \frac{F''(X_{Nf})}{2}x^2 + F'(X_{Nf})x + F(X_{Nf}). \quad (8)$$

If  $Z(x)$  and  $Z'(x)$  intersect the  $y$ -axis in points  $N_z$  and  $N_{z'}$  respectively, then (8) can be expressed as

$$Z(x) \equiv ax^4 - 6a\epsilon^2x^2 + y_{N_{z'}}x + y_{N_z} \quad (9)$$

where (see (4) and Figures 2, 3)

$$\begin{cases} \epsilon^2 = \frac{(3b^2 - 8ac)}{48a^2} \equiv \frac{-H}{48a^2} \equiv \frac{-p}{6a}, \\ y_{N_z} = F(X_{Nf}) \equiv \frac{I}{12a} - \frac{3H^2}{48^2a^3} \equiv r, \\ y_{N_{z'}} = F'(X_{Nf}) \equiv \frac{G}{8a^2} \equiv q, \\ -12a\epsilon^2 = F''(X_{Nf}). \end{cases} \quad (10)$$

Expressing the reduced quartic  $Z(x)$  in this form (Equation 9) greatly facilitates visualisation, since we can now 'see' how the configuration of the curves  $Z(x)$  and  $Z'(x)$  is related to the coefficients. For example (assuming  $a > 0$ ), if the  $x^2$  term is positive then  $\epsilon$  is complex ( $\epsilon^2 < 0$ ), and so the quartic will have two complex points of inflection and hence only one real turning point (cf. [10]).

If  $x_{T_i}$  are the  $x$ -coordinates of the turning points of  $Z(x)$ , then by differentiating (9) we have (see Equations 5a and 10)

$$4x_{T_1}x_{T_2}x_{T_3} = \frac{-y_{N_z'}}{a} = 8\sqrt{r_1r_2r_3}, \quad (11)$$

and hence the sign of  $\sqrt{r_1r_2r_3}$  is the same as that of  $-y_{N_z'}/a$  and  $x_{T_1}x_{T_2}x_{T_3}$ . It follows, therefore, that we can actually ‘see’ the correct sign of  $\sqrt{r_1r_2r_3}$  simply by observing the signs of the abscissae of the turning points of the reduced quartic, or by noting the location of  $N_z'$  in relation to the abscissa.

For example (assuming  $a > 0$ ), if the roots  $z_j$  are such that the middle turning point,  $T_2$ , is to the left of the  $y$ -axis, then not only will  $y_{N_z'}$  be negative (Figure 2) but just two of the three  $x_{T_i}$  will be negative resulting in a positive product for  $x_{T_1}x_{T_2}x_{T_3}$ , and hence  $\sqrt{r_1r_2r_3}$  will also be positive (see Equation 11). Conversely, if the middle turning point is to the right of the  $y$ -axis, then  $y_{N_z'}$  will be positive, and only one of the  $x_{T_i}$  will be negative making the product  $x_{T_1}x_{T_2}x_{T_3}$  negative.

## 5 Roots

As regards the roots  $z_j$  of the reduced quartic  $Z(x)$ , we can initially choose *any* sign combination for the  $\sqrt{r_i}$ , and then evaluate the sign of the product  $\sqrt{r_1r_2r_3}$ . If the sign of the product is the *same* as that of  $-y_{N_z'}/a$  (see Equation 11) then we have a valid combination of signs, and can proceed to determine the four  $z_j$  using (3). Otherwise, it is only necessary to change the sign of any *one* of the  $\sqrt{r_i}$  (say,  $\sqrt{r_1} \rightarrow -\sqrt{r_1}$ ), and proceed as before using (3).

When the reduced quartic is symmetric about the  $y$ -axis one of the  $x_{T_i}$  will be zero and hence the product  $\sqrt{r_1r_2r_3}$  is zero. However, the solution in this case is trivial since  $Z(x)$  is then an even function as  $y_{N_z'}$  is also zero.

## 6 Application

Since all resolvent cubics of the quartic can be transformed to a standard form [9], typically expressed as [6, p. 77]

$$T(x) \equiv x^3 - 3Ix + J, \quad (12)$$

we can solve any quartic by solving instead a simple reduced form of the resolvent, say  $T(x) = 0$ , and then recover the roots of Euler’s resolvent using the transformation which carries the reduced form back to  $R(x)$ .

For example, the translation  $x \mapsto x + x_{N_r}$  to reduce  $R(x)$ , for which  $x_{N_r} = -p/(6a) \equiv \varepsilon^2$ , generates the reduced form  $S(x)$ , as follows:

$$S(x) \equiv R(x + \varepsilon^2) \equiv x^3 - \frac{I}{48a^2}x + \frac{J}{1728a^3}. \quad (13)$$

The substitution  $x \mapsto x/(12a)$  then scales  $1728a^3S(x)$  to  $T(x)$ , and hence if the roots of  $S(x) = 0$  and  $T(x) = 0$  are  $s_i$  and  $t_i$  respectively, then

$$r_i = s_i + \varepsilon^2 = \frac{t_i}{12a} + \varepsilon^2. \quad (14)$$

This convenient approach is illustrated in Example 1.




Figure 3:

Euler’s resolvent cubic  $R(x)$  with three real roots ( $h_r^2 > y_{Nr}^2$ , i.e.  $4I^3 > J^2$ ) which are all positive ( $\epsilon^2 > 0$ ,  $x_{Nr}^2 > \delta_r^2$ ). The conditions  $\epsilon^2 > 0$ ,  $x_{Nr}^2 < \delta_r^2$  are associated with two negative roots (dashed curve). Note that  $G^2$ ,  $H$ ,  $I$ ,  $J$  are constant multiples respectively of the resolvent’s geometric parameters  $y_P$ ,  $x_{Nr}$ ,  $\delta_r^2$ ,  $y_{Nr}$  ( $\rho_1 = 64^2 a^6$ ,  $\rho_2 = 48a^2$ ,  $\rho_3 = 1728a^3$ ,  $\rho_4 = 12a$ ).

The invariants  $I$ ,  $J$  are readily visualised since any reduced cubic can be expressed in terms of its geometric parameters  $\delta$  and  $y_N$  as in [8]

$$Ax^3 - 3A\delta^2x + y_N = 0. \tag{15}$$

For example, equating coefficients between  $S(x)$ ,  $T(x)$  and the monic form of (15), and noting that  $h^2 = 4A^2\delta^6$  [8], shows that  $I$ ,  $J$  are simply constant multiples of  $\delta^2$ ,  $y_N$  as follows (Figure 3):

$$\begin{cases} A_r = A_s = A_t = 1, \\ I = \delta_r^2(12a)^2 = \delta_s^2(12a)^2 = \delta_t^2, \\ J = y_{Nr}(12a)^3 = y_{Ns}(12a)^3 = y_{Nt}, \\ \frac{4I^3}{J^2} = \left(\frac{h_r}{y_{Nr}}\right)^2 = \left(\frac{h_s}{y_{Ns}}\right)^2 = \left(\frac{h_t}{y_{Nt}}\right)^2. \end{cases} \tag{16}$$

Thus each of these invariants has a visible geometric interpretation in relation to Euler’s resolvent cubic, either as a position parameter with respect to the axes ( $G$ ,  $H$ ,  $J$ ), or as a shape parameter ( $I$ ). For example, we can now see that the condition  $J = 0$  simply indicates that the  $N$ -point of the resolvent cubic lies on the  $x$ -axis and all that that implies (see Example 2). Similarly, the condition  $I = 0$  indicates that the resolvent adopts the ‘cubic parabola’ form. Furthermore  $y_P \leq 0$ , which reveals how and why the resolvent cubic cannot have just a single negative root. The syzygy  $-27G^2 = H^3 - 48a^2IH + 64a^3J$  [6, p. 76] is generated by substituting into  $S(x)$  the coordinates of  $P$  ( $H/(48a^2)$ ,  $-G^2/(64^2a^6)$ ).

## 7 Euler's cubic and the quartic root configurations

A very significant but seemingly overlooked aspect of Euler's resolvent cubic is its beautiful and symmetric relationship with two important algebraic objects, namely the discriminant  $4I^3 - J^2$  and the seminvariant  $H^2 - 16a^2I$ , the signs of which distinguish between the various quartic root configurations [5, § 68; 6, p. 80; 7, p. 28]. Visualising the resolvent in relation to the invariants (Figure 3) reveals the mechanisms, as follows:

### 7.1 $4I^3 - J^2$

Since  $h^2 = 4A^2\delta^6$  [8], it follows from (16) that

$$\frac{-(4I^3 - J^2)}{12^6 a^6} = y_{Nr}^2 - h_r^2. \quad (17)$$

Thus the quartic discriminant  $4I^3 - J^2$  is simply a constant multiple of  $y_{Nr}^2 - h_r^2$ , the sign of which reflects whether the  $x$ -axis lies *between* ( $y_{Nr}^2 < h_r^2$ ), *on* ( $y_{Nr}^2 = h_r^2$ ), or *outside* ( $y_{Nr}^2 > h_r^2$ ) the turning points of the resolvent cubic (Figure 3).

### 7.2 $H^2 - 16a^2I$

The sign of this algebraic object distinguishes (when  $\varepsilon^2 > 0$ ) between the then two possible quartic root configurations associated with the case  $4I^3 - J^2 > 0$ , namely (a) four real roots ( $H^2 - 16a^2I > 0$ ), and (b) four complex roots ( $H^2 - 16a^2I < 0$ ) [5, § 68]. Substituting for  $H$  (Equation 10) and  $I$  (Equation 16) gives

$$H^2 - 16a^2I = (-48a^2\varepsilon^2)^2 - 16a^2(12^2 a^2 \delta_r^2) = 3^2 4^4 a^4 (\varepsilon^4 - \delta_r^2).$$

But  $\varepsilon^2 = x_{Nr}$  (Figure 3) and hence

$$\frac{H^2 - 16a^2I}{3^2 4^4 a^4} = x_{Nr}^2 - \delta_r^2. \quad (18)$$

Thus  $H^2 - 16a^2I$  is just a constant multiple of  $x_{Nr}^2 - \delta_r^2$ , the sign of which (when  $\varepsilon^2 > 0$ ) reflects whether the  $y$ -axis lies *between* ( $x_{Nr}^2 < \delta_r^2$ ), *on* ( $x_{Nr}^2 = \delta_r^2$ ), or *outside* ( $x_{Nr}^2 > \delta_r^2$ ) the turning points of the resolvent cubic (cf. [6, p. 80, proposition 7]).

For example (Figure 3), when a quartic with three real turning points ( $\varepsilon^2 > 0$ ) has four real roots ( $4I^3 - J^2 > 0$ ) Euler's cubic  $R(x)$  has three positive real roots—the  $y$ -axis lies *outside* the two turning points—and so  $x_{Nr}^2 > \delta_r^2$  and hence  $H^2 - 16a^2I > 0$ .

Conversely, when a quartic with three real turning points ( $\varepsilon^2 > 0$ ) has four complex roots ( $4I^3 - J^2 < 0$ ),  $R(x)$  then has exactly two negative real roots, and so its turning point  $T'$  (Figure 3) lies to the left of the  $y$ -axis ( $x_{Nr}^2 < \delta_r^2$ ), hence  $H^2 - 16a^2I < 0$ .

### 7.3 A new class of object?

Since  $H^2 - 16a^2I$  functions with regard to the  $y$ -axis in *exactly* the same way that  $4I^3 - J^2$  functions with regard to the  $x$ -axis, I would like to suggest that this pair of algebraic objects should be regarded as forming a distinct class of object—thereby linking two previously unrelated algebraic quantities with a single unifying concept.

## 8 Example 1

Solve  $F(X) \equiv X^4 - 11X^3 + 41X^2 - 61X + 30 = 0$ .

The key parameters are:  $a = 1$ ,  $X_{Nf} = 11/4$ ,  $Y_{Nf'} = F'(X_{Nf}) = -15/8$ ,  $I = 28$ ,  $J = -160$ ,  $\varepsilon^2 = 35/48$ . Using say,  $T(x)$ , we solve<sup>3</sup>

$$T(x) \equiv x^3 - 84x - 160 = 0,$$

the three  $t_i$  being  $-8, -2, 10$ . The  $\sqrt{r_i}$  are therefore given by

$$\begin{cases} \sqrt{r_1} = \sqrt{\varepsilon^2 + \frac{t_1}{12a}} = \sqrt{\frac{35}{48} - \frac{8}{12}} = \frac{1}{4}, \\ \sqrt{r_2} = \sqrt{\varepsilon^2 + \frac{t_2}{12a}} = \sqrt{\frac{35}{48} - \frac{2}{12}} = \frac{3}{4}, \\ \sqrt{r_3} = \sqrt{\varepsilon^2 + \frac{t_3}{12a}} = \sqrt{\frac{35}{48} + \frac{10}{12}} = \frac{5}{4}. \end{cases}$$

Since the sign of  $-Y_{Nf'}/a$  is positive then the product of the  $\sqrt{r_i}$  must also be positive—which it is. Finally, adding  $X_{Nf}$  recovers the quartic roots ( $X_j = X_{Nf} \pm \sqrt{r_1} \pm \sqrt{r_2} \pm \sqrt{r_3}$ ) using (3) as follows:

$$\begin{cases} X_1 = \frac{11}{4} - \frac{1}{4} + \frac{3}{4} - \frac{5}{4} = 2, \\ X_2 = \frac{11}{4} - \frac{1}{4} - \frac{3}{4} + \frac{5}{4} = 3, \\ X_3 = \frac{11}{4} + \frac{1}{4} + \frac{3}{4} + \frac{5}{4} = 5, \\ X_4 = \frac{11}{4} + \frac{1}{4} - \frac{3}{4} - \frac{5}{4} = 1. \end{cases}$$

Even the solution of  $T(x) = 0$  is greatly simplified since  $\delta$ ,  $h$ ,  $y_N$  are simple functions of  $I$  and  $J$  (see Equation 16). For example,  $T(x) = 0$  has three real roots in this case since  $(y_{Nt}/h_t)^2 \equiv J^2/(4I^3) \leq 1$  [8].

## 9 Example 2

*Explain the significance of  $J = 0$ ,  $I > 0$ , for a quartic with four real roots.*

The condition  $J = 0$  implies that Euler's resolvent cubic has its  $N$ -point on the  $x$ -axis (Figure 3), and hence it has three roots in arithmetic progression. If also  $I > 0$  (resolvent cubic has two real turning points), then the resolvent's roots are distinct and (with the root at infinity) form a harmonic range. Since the roots of the parent quartic have the same cross-ratio they also form a harmonic range.

<sup>3</sup>Note that we could instead solve  $S(x) = 0$ , and then use  $r_i = \varepsilon^2 + s_i$  (see Equation 14).

## 10 Acknowledgements

I would like to thank Professor JE Cremona (University of Warwick) for reading drafts of this paper and for his constructive comments. I also acknowledge the helpful suggestions of the anonymous reviewer.

## 11 References

- 1 L. Euler. De formis radicum aequationum cujusque ordinis conjectatio. *Commentarii academiae scientiarum imperialis Petropolitanae* (1733); **6** (pub. 1738), pp. 216–231 = *Opera Omnia*, Series 1 (Pure mathematics), **6** (Theory of equations), 1–19. [Euler Archive, E30 (Latin): <http://math.dartmouth.edu/~euler/pages/E030.html>]
- 2 Bell J (2008). A conjecture on the forms of the roots of equations. *arXiv:0806.1927v1* [math.HO]. <http://arxiv.org/abs/0806.1927> [An English translation of Euler’s *De formis radicum aequationum cujusque ordinis conjectatio* (E30)].
- 3 L. Euler. *Vollständige Anleitung zur Algebra* (Elements of algebra), 2 vols, Royal Academy of Sciences, St. Petersburg (1770). [Euler Archive, E387 (English): <http://math.dartmouth.edu/~euler/docs/originals/E387e.P1S4.pdf>]
- 4 C. R. Sangwin (Ed.). Euler’s *Elements of Algebra*. Tarquin Publications, St Albans, UK (2006). [English translation of Euler 1770 (E387)]
- 5 W. S. Burnside and A. W. Panton. *The theory of equations: with an introduction to the theory of binary algebraic forms*. (7th edn.) 2 vols; Longmans, Green and Co., London (1912).
- 6 J. E. Cremona. Reduction of binary cubic and quartic forms. *J. Comput. Math.*, **2** (1999), pp. 62–92. <http://www.lms.ac.uk/jcm/2/lms98007/> [The seminvariants  $G$  and  $H^2 - 16a^2I$  are denoted here by  $R$  and  $3Q$  respectively]
- 7 P. J. Olver. *Classical invariant theory*. London Mathematical Society Student Texts No. 44. Cambridge University Press (1999).
- 8 R. W. D. Nickalls. A new approach to solving the cubic: Cardan’s solution revealed. *Math. Gaz.*, **77** (Nov 1993) pp. 354–359. <http://www.nickalls.org/dick/papers/math/cubic1993.pdf>. <http://www.jstor.org/stable/3619777>
- 9 R. S. Ball. Note on the algebraical solution of biquadratic equations. *Quarterly Journal of Pure and Applied Mathematics*, **7** (1866) pp. 6–9, 358–369. <http://www.nickalls.org/dick/papers/math/ball1866quartic.pdf>.
- 10 J. P. Dalton. On the graphical discrimination of the cubic and of the quartic. *Math. Gaz.*, **17** (July 1933) pp. 189–196 <http://www.jstor.org/stable/3607613>

◇ ◇ ◇